



View this article online at: patient.info/doctor/generalised-anxiety-disorder-assessment-gad-7

Generalised Anxiety Disorder Assessment (GAD-7)

This easy-to-use self-administered patient questionnaire is used as a screening tool and severity measure for generalised anxiety disorder (GAD).^{[1] [2]}

Generalised Anxiety Disorder Questionnaire (GAD-7)

Over the last 2 weeks, how often have you been bothered by any of the following problems?

Feeling nervous, anxious or on edge?

Not at all
Several days
More than half the days
Nearly every day

Not being able to stop or control worrying?

Not at all
Several days
More than half the days
Nearly every day

Worrying too much about different things?

Not at all
Several days
More than half the days
Nearly every day

Trouble relaxing?

Not at all
Several days
More than half the days
Nearly every day

Being so restless that it is hard to sit still?

Not at all
Several days
More than half the days
Nearly every day

Becoming easily annoyed or irritable?

Not at all
Several days
More than half the days
Nearly every day

Feeling afraid as if something awful might happen?

Not at all
Several days
More than half the days
Nearly every day

Total= /21

The GAD-7 originates from Spitzer RL, Kroenke K, Williams JB, et al; A brief measure for assessing generalized anxiety disorder: the GAD-7. Arch Intern Med. 2006 May 22;166(10):1092-7. GAD-7 © Pfizer Inc. all rights reserved; used with permission.

The GAD-7 score is calculated by assigning scores of 0, 1, 2, and 3, to the response categories of 'not at all', 'several days', 'more than half the days', and 'nearly every day', respectively, and adding together the scores for the seven questions.

Scores of 5, 10, and 15 are taken as the cut-off points for mild, moderate and severe anxiety, respectively. When used as a screening tool, further evaluation is recommended when the score is 10 or greater.

Using the threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD. It is moderately good at screening three other common anxiety disorders - panic disorder (sensitivity 74%, specificity 81%), social anxiety disorder (sensitivity 72%, specificity 80%) and post-traumatic stress disorder (sensitivity 66%, specificity 81%).^[3]

Further reading & references

1. Swinson RP; The GAD-7 scale was accurate for diagnosing generalised anxiety disorder. Evid Based Med. 2006 Dec;11(6):184.
2. Spitzer RL, Kroenke K, Williams JB, et al; A brief measure for assessing generalized anxiety disorder: the GAD-7. Arch Intern Med. 2006 May 22;166(10):1092-7.
3. Kroenke K, Spitzer RL, Williams JB, et al; Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. Ann Intern Med. 2007 Mar 6;146(5):317-25.





Disclaimer: This article is for information only and should not be used for the diagnosis or treatment of medical conditions. EMIS has used all reasonable care in compiling the information but make no warranty as to its accuracy. Consult a doctor or other health care professional for diagnosis and treatment of medical conditions. For details see our [conditions](#).

Original Author: Dr Huw Thomas	Current Version: Dr Hayley Willacy	Peer Reviewer: Dr Colin Tidy
Document ID: 8736 (v11)	Last Checked: 28/01/2016	Next Review: 26/01/2021

View this article online at: patient.info/doctor/generalised-anxiety-disorder-assessment-gad-7

Discuss Generalised Anxiety Disorder Assessment (GAD-7) and find more trusted resources at [Patient](#).

Ask your doctor about Patient Access

-  Book appointments
-  Order repeat prescriptions
-  View your medical record
-  Create a personal health record (iOS only)



Simple, quick and convenient.
Visit patient.info/patient-access
or search 'Patient Access'

Like us.
it's good for you!

 Patient

Like us on **facebook**
fb.com/patient

